

Short notes

Karyotypes of six previously unstudied European mealybugs (Homoptera: Pseudococcidae)

I.A. Gavrilov, I.V. Trapeznikova

Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St. Petersburg, 199034, Russia.

E-mail: coccids@gmail.com

Abstract. *Phenacoccus peruvianus* Granada de Willink, 2007, *Ph. prope avenae* Borchsenius, 1949, *Ph. hordei* (Lindeman, 1886), *Balanococcus boratynskii* Williams, 1962, *Trionymus radicum* (Newstead, 1895), *Rhizoecus halophilus* (Hardy, 1868) were studied karyologically for the first time. All species demonstrate $2n=10$ and a Lecanoid genetic system. Photos of karyotypes of all studied species are given.

Key words: mealybugs, scale insects, karyotype, chromosome number.

By now, only 117 species of mealybugs, that is about 5 % of the world fauna of the family Pseudococcidae, have been studied cytogenetically (see catalogue of Gavrilov, 2007; Gavrilov, Trapeznikova, 2007). Here, we present the karyotypes of 6 further mealybug species, collected in recent years in Portugal, Bulgaria and Russia. All species demonstrate a Lecanoid genetic system characteristic of most Pseudococcidae species (Nur, 1980; Nur et al., 1987; Gavrilov, 2007). Once males display this system, they are diploid, one haploid set of chromosomes being heterochromatinized in most cells (see Figs 2, 5, 7).

Chromosomal preparations were performed as previously described (Gavrilov, Trapeznikova, 2007, 2008). All material is

deposited at the Zoological Institute, Russian Academy of Sciences.

Phenacoccus peruvianus Granada de Willink, 2007 (Fig. 1)


Material. K № 550, Portugal, Lisboa vicinity, *Pittosporum* sp., 26.IX.2007, I. Gavrilov.

Embryos from female body. $2n=10$.

Phenacoccus prope avenae Borchsenius, 1949 (Fig. 2)

Material. K № 555, Portugal, Lisboa vicinity, undetermined bush, 27.IX.2007, I. Gavrilov.

Embryos from female body. $2n=10$.



Figs 1-8. Karyotypes of six mealybug species. **1** - *Phenacoccus peruvianus*, female embryonic cell. **2** - *Ph. prope avenae*, female embryonic cell. **3** - *Ph. hordei*, female embryonic cell. **4-5** - *Balanococcus boratynskii*; **4** - female embryonic cell, **5** - male embryonic cell. **6-7** - *Trionymus radicum*; **6** - female embryonic cell, **7** - male embryonic cell. **8** - *Rhizoecus halophilus*, female embryonic cell. Arrows indicate a heterochromatinized haploid set of chromosomes. Bar = 10 μ m.

Phenacoccus hordei (Lindeman, 1886) (Fig. 3)

Material. M № 1, Russia, Belgorod province, Gubkin, under the leaf sheaths of *Poa* sp., 1.VI.2007, I. Trapeznikova. M № 2, Belgorod province, Valuiki, 2.VI.2007. M № 9, the same, but Bir'uch, 14.VI.2007. M № 13, Belgorod province, 30 km S Bir'uch, 6.VII.2007. M № 30, Belgorod province,

Novii Oskol, 8.VII.2008. M № 31, Belgorod province, Alekseevka, 9.VII.2008.

Embryos from the female body. $2n=10$.

Balanococcus boratynskii Williams, 1962 (Figs 4-5)

Material. K № 572, Bulgaria, Vitosha mt., alpine zone, under the leaf sheath of undetermined grass, 6.VII.2008, I. Gavrilov.

Embryos from the female body. $2n=10$.

Comments. The second cytogenetically studied species of the genus, after *B. singularis* (Schmutterer, 1952) (Gavrilov, 2007; Gavrilov, Trapeznikova, 2007).

Trionymus radicum (Newstead, 1895)
(Figs 6-7)

Material. K № 573, 574, 575, Bulgaria, Rila mts., Yastrebits, 2300 m, under the leaf sheaths of undetermined grass, 09.VII.2008, I. Gavrilov. K № 576, Bulgaria, Rila mts., Yastrebits, 2300 m, on stems of *Dianthus* sp., 9.VII.2008, I. Gavrilov.

Embryos from female body. $2n=10$.

Comments. The genus *Trionymus* Berg, 1899 provides a good possibility for karyotaxonomic conclusions. Gavrilov (2004, 2007) and Gavrilov and Trapeznikova, 2007 have shown that the type species of the genus, *T. perrisii* (Signoret, 1875), and also *T. aberrans* Goux, 1938 and *T. haancheni* (McKenzie, 1960) have $2n=16$ and bear similarity in morphology. On the other hand, the species that are not very similar in morphology to the type species, demonstrate different chromosome numbers. For example, *T. multivorus* (Kiritshenko, 1936) having $2n=10$ was transferred to the genus *Dysmicoccus* Ferris, 1950 (Gavrilov, 2004). Similarly, *T. radicum* has $2n=10$ and tubular ducts with collars inherent in the genus *Balanococcus* Williams, 1962 arguing for the transferring this species to the genus *Balanococcus*.

Rhizoecus halophilus (Hardy, 1868)
(Fig. 8)

Material. K № 578, Bulgaria, Rila mts., the road to Musalka mt. near Musala lakes, ≈ 2300 m, on thin roots of undetermined grass, 11.VII.2008, I. Gavrilov.

Embryos from ovisacs. $2n=10$.

Comments. To date, the karyotypes of 4 species of the genus *Rhizoecus* Künckel

d'Herculais, 1878, with chromosome numbers $2n=8$, $2n=10$ and $2n=12$, have been reported (McKenzie, 1967; Nur et al., 1987; Gavrilov, 2004). *Rh. halophilus*, studied here, has the same chromosome number as in *Rh. mayanus* (Hambleton, 1946).

ACKNOWLEDGEMENTS

The work was supported by the Russian Foundation for Basic Researches (projects № 08-04-00787 and № 09-04-91229-Cra), by the grant of the President of Russian Federation (MK-6075.2010.4) and special grant of the St. Petersburg Government.

REFERENCES

- Gavrilov I.A. 2004. Taxonomic and cytogenetic studies of scale insects (Homoptera: Coccinea) of European Russia // *Proc. Zool. Inst. RAS*. 300: 77-82.
- Gavrilov I.A. 2007. A catalogue of chromosomal numbers and genetic systems of scale insects (Homoptera: Coccinea) of the world // *Israel J. Entomol.* 37: 1-53.
- Gavrilov I.A., Trapeznikova I.V. 2007. Karyotypes and reproductive biology of some mealybugs (Insecta: Coccinea: Pseudococcidae) // *Comp. Cytogenet.* 1(2): 139-148.
- Gavrilov I.A., Trapeznikova I.V. 2008. Cytogenetic studies of European Pulvinariini (Homoptera: Coccidae) // *Comp. Cytogenet.* 2008(2): 131-138.
- McKenzie H.L. 1967. Mealybugs of California with taxonomy, biology and control of North American species (Homoptera: Coccoidea: Pseudococcidae). Berkeley, Los Angeles. 525 p.
- Nur U. 1980. Evolution of unusual chromosome systems in scale insects (Coccoidea: Homoptera) (pp. 97-117) // Blackman R.L., Hewitt G.M., Ashburner M. (Eds). *Insect Cytogenetics*. London. 278 p.
- Nur U., Brown S.W., Beardsley J.W. 1987. Evolution of chromosome number in mealybugs (Pseudococcidae: Homoptera) // *Genetica*. 74: 53-60.

Received October 29, 2010.

Accepted by V.G. Kuznetsova, December 14, 2010.

Published December 30, 2010.